



Calhoun: The NPS Institutional Archive
DSpace Repository

Faculty and Researchers

Faculty and Researchers' Publications

1977-03

A note on a confidence interval for an interclass mean

Jayachandran, T.

Wiley Online

Jayachandran, T. "A note on a confidence interval for an interclass mean." Naval Research Logistics Quarterly 24.1 (1977): 197-200.
<http://hdl.handle.net/10945/60197>

This publication is a work of the U.S. Government as defined in Title 17, United States Code, Section 101. Copyright protection is not available for this work in the United States.

Downloaded from NPS Archive: Calhoun



Calhoun is the Naval Postgraduate School's public access digital repository for research materials and institutional publications created by the NPS community. Calhoun is named for Professor of Mathematics Guy K. Calhoun, NPS's first appointed -- and published -- scholarly author.

Dudley Knox Library / Naval Postgraduate School
411 Dyer Road / 1 University Circle
Monterey, California USA 93943

<http://www.nps.edu/library>

A NOTE ON A CONFIDENCE INTERVAL FOR AN INTERCLASS MEAN

T. Jayachandran*

*Naval Postgraduate School
Monterey, California*

ABSTRACT

An exact confidence interval for an interclass mean, that is, the mean of a composite sample made up several subsamples of unequal sizes n_i is presented.

1. INTRODUCTION

Suppose $X_{ij}, j=1, 2, \dots, n_i; i=1, 2, \dots, k$ is a composite sample of size

$$N = \sum_{i=1}^k n_i$$

that is comprised of k subsamples of sizes n_i . The i^{th} subsample is a random sample from a normal distribution $N(a_i, \sigma^2)$ and the a_i 's, $i=1, 2, \dots, k$ are assumed to be independent and identically distributed (*i.i.d.*) as $N(\mu, \sigma_a^2)$. μ is sometimes known as the interclass mean and the problem considered in this note is the construction of a confidence interval for μ . Some practical uses of such an interval are discussed in a paper by Long [1]. For example, the composite sample may be measurements on a characteristic of the output of a factory made on different days. μ would correspond to the true mean value of the characteristic being studied.

The construction of a confidence interval for μ is straightforward if all the n_i are equal. For unequal subsample sizes Long [1] obtained an approximate interval that he shows to be reasonably accurate. The procedure proposed in this paper leads to an exact confidence interval for μ in the case of unequal n_i .

2. PRELIMINARIES

The observations $X_{ij}, j=1, 2, \dots, n_i; i=1, 2, \dots, k$ may be assumed to satisfy the variance components model

$$(2.1) \quad Y_{ij} = a_i + e_{ij}, j=1, 2, \dots, n_i; i=1, 2, \dots, k$$

*This research was supported by the Office of Naval Research as part of the Foundation Research Program at the Naval Postgraduate School.

where the a_i 's are *i.i.d.* $N(\mu\sigma_a^2)$, the e_{ij} 's are *i.i.d.* $N(0, \sigma^2)$ and the a_i 's and e_{ij} 's are mutually independent. Let

$$\bar{X}_i = \sum_{j=1}^{n_i} \frac{X_{ij}}{n_i} \text{ and } \bar{X} = \sum_{i=1}^k \frac{\bar{X}_i}{k}$$

and

$$S^2 = \sum_{i=1}^k \frac{(\bar{X}_i - \bar{X})^2}{k-1}$$

Then, \bar{X}_i , $i=1, 2, \dots, k$ are independent and normally distributed

$$N\left(\mu, \sigma_a^2 + \frac{\sigma^2}{n_i}\right).$$

If $n_i=n$ for all i the \bar{X}_i will constitute a random sample from a normal distribution

$$N\left(\mu, \sigma_a^2 + \frac{\sigma^2}{n}\right).$$

Thus,

$$(2.2) \quad T = \frac{k^{1/2}(\bar{X} - \mu)}{S}$$

has a student's t distribution with $(k-1)$ degrees of freedom, and this leads to a confidence interval for μ .

For the case where the n_i are not all equal, Long [1] obtained an approximate confidence interval for μ by treating T as a student's t variable. On the basis of a study of the exact distribution of T for $k=2, 3$ and an examination of the moments of T for large values of k , Long [1] concludes that the t approximation is fairly reliable. He also points out that the approximation is likely to go wrong if there are wide variations in the subsample sizes n_i or if σ^2 is large relative to σ_a^2 .

3. EXACT CONFIDENCE INTERVAL WITH UNEQUAL SUBSAMPLE SIZES

For unequal subsample sizes, the \bar{X}_i are not identically distributed since the variance of \bar{X}_i is

$$\sigma_a^2 + \frac{\sigma^2}{n_i},$$

even though all of them have the same mean μ . Thus, the construction of a student's t variable, based on \bar{X}_i only, independent of the nuisance parameters $\sigma_a^2 + \sigma^2/n_i$, is not possible. To get around this difficulty let

$$(3.1) \quad Z_i = C_{ii}X_{ii} + C_{i2}\bar{X}_i \quad i=1, 2, \dots, k$$

where

$$C_{ii} = \left(\frac{n_i C - 1}{n_i - 1} \right)^{1/2}$$

$$C_{i2} = 1 - C_{ii}$$

and

$$C = \frac{1}{\min n_i}.$$

Then, as shown in the appendix,

$$E(Z_i) = \mu$$

$$V(Z_i) = \sigma_a^2 + C\sigma^2 = \sigma_a^2 + \frac{\sigma^2}{\min n_i}$$

If

$$\bar{Z} = \frac{1}{k} \sum_{i=1}^k Z_i$$

and

$$S_z^2 = \frac{1}{k-1} \sum_{i=1}^k (Z_i - \bar{Z})^2$$

then

$$(3.2) \quad T = \frac{k^{1/2}(\bar{Z} - \mu)}{S_z}$$

has a student's t -distribution with $k-1$ degrees of freedom. An exact confidence interval for μ is now obtainable in the usual way.

4. DISCUSSION

The exact procedure proposed in Section 3 is somewhat ad hoc in nature in the sense that the first observation X_{i1} in each subsample plays a prominent role. To avoid any systematic bias that may creep in, it would be preferable to randomly permute the observations in each of the k subsamples before applying the procedure. If all the subsample sizes n_i are equal to n , the statistic (3.2) reduces to the usual T statistic (2.2).

5. APPENDIX

Under the assumptions of the variance components model (2.1)

$$E(X_{ij}) = \mu \quad j=1, 2, \dots, n_i; i=1, 2, \dots, k$$

$$\text{Var}(X_{ij}) = \sigma^2 + \sigma_a^2$$

and

$$\text{Cov}(X_{ij}, X_{i'j'}) = \begin{cases} 0 & i \neq i' \\ \sigma_a^2 & i = i', j \neq j' \end{cases}$$

It follows

$$E(\bar{X}_i) = \mu$$

$$V(\bar{X}_i) = \sigma_a^2 + \frac{\sigma^2}{n_i} \quad i=1, 2, \dots, k$$

and

$$\text{Cov}(X_{ij}, \bar{X}_i) = \sigma_a^2 + \frac{\sigma^2}{n_i} \quad j=1, 2, \dots, n_i$$

Therefore, if

$$Z_i = C_{i1}X_{i1} + C_{i2}\bar{X}_i$$

then

$$E(Z_i) = (C_{i1} + C_{i2})\mu = \mu$$

and

$$V(Z_i) = C_{i1}^2 V(X_{i1}) + C_{i2}^2 V(\bar{X}_i) + 2C_{i1}C_{i2} \text{Cov}(X_{i1}, \bar{X}_i)$$

$$= C_{i1}^2(\sigma^2 + \sigma_a^2) + (C_{i2}^2 + 2C_{i1}C_{i2})\left(\sigma_a^2 + \frac{\sigma^2}{n_i}\right).$$

substitution of

$$C_{i1} = \left(\frac{n_i C - 1}{n_i - 1}\right)^{1/2}, \quad C_{i2} = 1 - C_{i1} \quad \text{and} \quad C = \frac{1}{\min n_i}$$

results in

$$E(Z_i) = \mu \quad \text{and} \quad V(Z_i) = \sigma_a^2 + \frac{\sigma^2}{\min n_i}.$$

REFERENCES

1. Long, W. M., "Estimation Problems When a Simple Type of Heterogeneity is Present in the Sample," *Biometrika*, 38, 90-101 (1951).